

Google scholar

function * "Common gamma chain" * structure

Search

Advanced Scholar Search

Scholar Preferences

Scholar Articles and patents

anytime

include citations

☒ Create email alert

Results 1 - 10 of about 1,860. (0.13 sec)

The structure and function of gamma c-dependent cytokines and receptors: regulation of T lymphocyte development and homeostasis.

TW He. TR Malek - Critical reviews in immunology, 1998 - ncbi.nlm.nih.gov

1: Crit Rev Immunol. 1998;18(6):503-24. The structure and function of gamma c-dependent cytokines and receptors ... Five cytokines, IL-2, IL-4, IL-7, IL-9, and IL-15, form one group that is characterized by utilizing the common gamma chain (gamma c) as a receptor subunit ...

Cited by 61 - Related articles - BL Direct - All 6 versions

Jak3-independent trafficking of the common (gamma) chain receptor subunit: chaperone function of Jaks revisited

nib.gov [HTML]

SR Holmann, AG Lam, S Frank, YJ Zhou, ... - and cellular biology, 2004 - Am Soc Microbiol ... with c in living cells appears to be a very sensitive indicator of proper Jak structure. ... regulation of c and Jak3 will be important given the critical in vivo functions of both ... The amino terminus of JAK3 is necessary and sufficient for binding to the common gamma chain and confers ...

Cited by 28 - Related articles - BL Direct - All 6 versions

... IL-13 bind to a shared heterodimeric complex on endothelial cells mediating vascular cell adhesion molecule-1 induction in the absence of the common gamma chain

hematologylibrary.org [PDF]

B Schnyder, S Lugli, N Faug, H Elber, RA ... - 1996 - bloodjournal.hematologylibrary.org ... Therefore, the subunit structure of the functional receptors for IL-4 and IL-13 on human endothelial cells does not use or require the common gamma chain of the IL ... with the 130-kD IL4R and enhances its affinity with the ligand. IL-13 and IL-4 have remarkably similar functions on dif ...

Cited by 49 - Related articles - BL Direct - All 6 versions

Blocking the common (gamma)-chain of cytokine receptors induces T cell apoptosis and long-term islet allograft survival

immunol.org [HTML]

XC Li, A Ima, Y Li, XX Zheng, TR Malek, ... - The Journal of ... 2000 - Am Assoc Immunol ... complex can bind IL-2 but cannot initiate signaling events, and, therefore, this structure is often ... Expression and function of the subunit of the IL-2, IL-4, and IL-7 ... Strom, and XC Li Islet Allograft Rejection in Nonobese Diabetic Mice Involves the Common (gamma)-Chain and CD28 ...

Cited by 89 - Related articles - BL Direct - All 6 versions

The common (gamma)-chain for multiple cytokine receptors

K Sugamura, H Asao, M Kuroki, N Tanaka, N ... - Advances in ... 1995 - Elsevier

... Structure and Function of IL-2 Receptor Subunits 227 111. Sharing of IL-2Ry among Multiple Cytokine Receptors 238 14. Causative Relationship between IL-2Ry Mutations and Human XSCID 246 V. Signal Transduction Initiated by the Receptors Sharing the gamma-Chain 253 Vi ...

Cited by 143 - Related articles - BL Direct - All 6 versions

Molecular cloning of a novel type 1 cytokine receptor similar to the common gamma chain

hematologylibrary.org [HTML]

K Fujio, T Iwakawa, T Kojima, T ... - 2000 - bloodjournal.hematologylibrary.org

... Molecular cloning of a novel type 1 cytokine receptor similar to the common gamma chain. ... View larger version (3622K): [in this window] [in a new window]. Fig 4. Structure of chimeric ... We next examined the function of the cytoplasmic domain of 1; chimeric receptors between 1 ...

Cited by 39 - Related articles - BL Direct - All 6 versions

Structure, function, and targeting of interleukin 4 receptors on human head and neck cancer cells

ascrjournals.org [HTML]

K Kawakami, P Leland, FK Pun - Cancer research, 2000 - AACR

... Experimental Therapeutics. Structure, Function, and Targeting of Interleukin 4 Receptors on Human Head and Neck Cancer Cells. ... However, IL-4R structure, function, and cytotoxic activity of IL-4 toxin in SCCHN have not been investigated. ...

Cited by 37 - Related articles - BL Direct - All 6 versions

Structure of the Quaternary Complex of Interleukin-2 with Its (alpha),(beta), and (gamma) c Receptors

X Wang, M Rickert, KC Garcia - Science's STKE, 2005 - www.scripps.org

... that acts through a quaternary receptor signaling complex containing alpha (IL-2R) beta (IL-2R), and common gamma chain (gamma c). IL-4, IL-7, IL-9, IL-15, and IL-21 c forms degenerate contacts with IL-2. The structure of c provides a rationale for loss-of-function mutations found ...

Cited by 79 - Related articles - All 13 versions

Stable and functional lymphoid reconstitution of common cytokine receptor gamma chain deficient mice by retroviral-mediated gene transfer

hematologylibrary.org [HTML]

C Schneider, T Shih, LI Shearer, D Guy, ... - 2004 - bloodjournal.hematologylibrary.org

... Phenotype and function of peripheral lymphocytes from c-transduced mice. ... B-cell responses in c-transduced animals. To assess B-cell function in c-transduced mice, we first analyzed steady-state circulating plasma immunoglobulin levels in animals >8 weeks posttransplant. ...

Cited by 145 - Related articles - BL Direct - All 6 versions


The interleukin-4 receptor activates STAT5 by a mechanism that relies upon common gamma-chain

jbc.org [HTML]

A Lucifora, P Mongelli, S Brandwein, S Berchtold, ... - Journal of Biological ... 1998 - ASBMB ... substrate), phosphatidylinositol 3-kinase, STAT6, and probably other proteins generate a network

of protein-protein contacts mediated by interactions between phosphotyrosines and Src homology
2 domains (9). Despite **structure-function** studies by several laboratories (10-16 ...

[Read for 5 min](#) - [Related articles](#) - [Bib. Download](#) - [All 6 versions](#)

 [Create email alert](#)

Google

Result Page: 1 2 3 4 5 6 7 8 9 10 [Next](#)

function * "Common gamma chain" *

[Go to Google Home](#) - [About Google](#) - [About Google Scholar](#)

©2010 Google